arXiv:1410.7898v2 [math.NT] 12 May 2015

ARITHMETIC PROPERTIES OF OVERPARTITION TRIPLES

LIUQUAN WANG

ABSTRACT. Let $\overline{p}_3(n)$ be the number of overpartition triples of n. By elementary series manipulations, we establish some congruences for $\overline{p}_3(n)$ modulo small powers of 2, such as

$$\overline{p}_3(16n+14)\equiv 0 \pmod{32}, \quad \overline{p}_3(8n+7)\equiv 0 \pmod{64}.$$

We also find many arithmetic properties for $\overline{p}_3(n)$ modulo 7, 9 and 11, involving the following infinite families of Ramanujan-type congruences: for any integers $\alpha\geq 1$ and $n\geq 0$, we have $\overline{p}_3\big(3^{2\alpha+1}(3n+2)\big)\equiv 0 \pmod{9\cdot 2^4}$, $\overline{p}_3(4^{\alpha-1}(56n+49))\equiv 0 \pmod 7$ and

$$\overline{p}_3\big(7^{2\alpha+1}(7n+3)\big)\equiv \overline{p}_3\big(7^{2\alpha+1}(7n+5)\big)\equiv \overline{p}_3\big(7^{2\alpha+1}(7n+6)\big)\equiv 0 \pmod 7.$$

1. INTRODUCTION

An overpartition ω of n is a partition of n in which the first occurrence of a part may be overlined. An overpartition k-tuple $(\omega_1,\omega_2,\cdots,\omega_k)$ of n is a k-tuple of overpartitions where the sum of all of the parts is n. We denote by $\overline{p}_k(n)$ the number of overpartition k-tuples of a positive integer n and we define $\overline{p}_k(0)=1$ for convenience. In particular, for $k=1$, $\overline{p}_1(n)$ is denoted as $\overline{p}(n)$. For $k=2$, $\overline{p}_2(n)$ is often denoted as $\overline{pp}(n)$.

We recall that two of Ramanujan's theta functions are

$$\varphi(q)=\sum_{n=-\infty}^{\infty}q^{n^2}=\frac{(-q;-q)_\infty}{(q;-q)_\infty}=\frac{(-q;-q)_\infty^2}{(q^2;q^2)_\infty},$$

$$\psi(q)=\sum_{n=0}^{\infty}q^{n(n+1)/2}=\frac{(q^2;q^2)_\infty^2}{(q;q)_\infty},$$

where $(a;q)_\infty=\prod_{n\geq 0}(1-aq^n)$ is standard q series notation. It is not difficult to find that the generating function of $\overline{p}_k(n)$ is

$$\sum_{n\geq 0}\overline{p}_k(n)q^n=\Big(\frac{(-q;q)_\infty}{(q;q)_\infty}\Big)^k=\frac{1}{\varphi(-q)^k}.$$

Arithmetic properties of $\overline{p}(n)$ and $\overline{pp}(n)$ have been investigated by many mathematicians. Numerous results have been obtained, see [4]–[10] and [12]-[21]. Here we shall mention some research which motivates this work.

By using 2-adic expansions of $\overline{p}(n)$ or the theory of modular forms, people found many congruences for $\overline{p}(n)$ modulo powers of 2. For example, Mahlburg [20] showed that $\overline{p}(n)\equiv 0 \pmod{64}$ holds for a set of integers of arithmetic density 1. Kim [15]

2010 *Mathematics Subject Classification.* Primary 05A17; Secondary 11P83.

Key words and phrases. Partitions; overpartition triples; congruences; theta functions.

gave the modulo 8 determination of $\overline{p}(n)$. He [14] also proved that $\overline{p}(n)$ is divisible by 128 for almost all positive integers. Similar results for $\overline{pp}(n)$ have also been found, see [16] for details. There are also some interesting Ramanujan-type congruences modulo small powers of 2. For example, Chen and Lin [6] proved that

$$\overline{pp}(8n+7) \equiv 0 \pmod{64}.$$

Keister, Sellers and Vary [18] showed that if $k = 2^m r$ where $m \geq 0$, and $r \geq 1$ is an odd integer, then

$$\overline{p}_k(n) \equiv \begin{cases} 2^{m+1} & \text{if } n \text{ is a square or twice a square} \\ 0 & \text{otherwise} \end{cases} \pmod{2^{m+2}}. \tag{1.1}$$

Following their steps, in Section 2, we give the modulo 16 determination of $\overline{p}_3(n)$. Meanwhile, we establish some congruences for $\overline{p}_3(n)$ modulo small powers of 2. For example, we have

$$\overline{p}_3(16n+14) \equiv 0 \pmod{32}, \quad \overline{p}_3(8n+7) \equiv 0 \pmod{64}.$$

In 2005, Hirschhorn and Sellers [12] obtained an infinite family of congruences for $\overline{p}(n)$ modulo 3: for any nonnegative integers n and α,

$$\overline{p}(9^\alpha(27n+18)) \equiv 0 \pmod{3}.$$

Chen and Lin [6] gave two infinite families of congruences for $\overline{pp}(n)$ modulo 3: for any integers $\alpha \geq 1$ and $n \geq 0$,

$$\overline{pp}(9^\alpha(3n+1)) \equiv \overline{pp}(9^\alpha(3n+2)) \equiv 0 \pmod{3}.$$

Inspired by their work, in Section 3, we will present an infinite family of congruences for $\overline{p}_3(n)$ modulo 9. For any integers $\alpha \geq 1$ and $n \geq 0$, we will show that

$$\overline{p}_3(3^{2\alpha+1}(3n+2)) \equiv 0 \pmod{9 \cdot 2^4}.$$

It seems much harder to obtain congruences for $\overline{p}_k(n)$ modulo primes other than 2 and 3. The methods used in this direction are quite different. We mention two results here. By using (p,k)-parametrization of theta functions, Chen and Xia [8] proved a conjecture of Hirschhorn and Sellers [13]:

$$\overline{p}(40n+35) \equiv 0 \pmod{40}.$$

Their proof is relatively long and two different short proofs of this conjecture were then given by Lin [17] and the author [21].

In 2012, Chen and Lin [6] found two infinite families of congruences satisfied by $\overline{pp}(n)$. For any integers $\alpha \geq 1$ and $n \geq 0$, we have

$$\overline{pp}(5^\alpha(5n+2)) \equiv \overline{pp}(5^\alpha(5n+3)) \equiv 0 \pmod{5}.$$

For nonnegative integers n and k, let $r_k(n)$ denote the number of representations of n as sum of k squares. In Section 4, we establish the following two arithmetic relations: For any integer $n \geq 1$,

$$\overline{p}_3(7n) \equiv (-1)^n r_3(n) \pmod{7},$$

and

$$\overline{p}_3(11n) \equiv (-1)^n r_7(n) \pmod{11}.$$

From these two relations we deduce many interesting congruences such as $\overline{p}_3(4^{\alpha-1}(56n+49)) \equiv 0 \pmod 7$ and

$$\overline{p}_3\big(7^{2\alpha+1}(7n+3)\big) \equiv \overline{p}_3\big(7^{2\alpha+1}(7n+5)\big) \equiv \overline{p}_3\big(7^{2\alpha+1}(7n+6)\big) \equiv 0 \pmod 7.$$

2. Congruences Modulo Powers of 2

Theorem 2.1. *We have*

$$\overline{p}_k(n) \equiv \begin{cases} -2k, & \text{if } n \text{ is an even square;} \\ 2k, & \text{if } n \text{ is an odd square;} \\ 2k(k+1), & \text{if } n \text{ is twice a square;} \\ 0, & \text{else} \end{cases} \pmod{2^{m(k)}}.$$

where $m(k) = 4$ when $k \equiv 0, 3 \pmod 4$ and $m(k) = 3$ otherwise .

Comparing with (1.1), it is easy to see that this result has the advantage when k is odd.

To prove this theorem, we need a simple lemma as follows.

Lemma 2.1. *Let $\chi(n)$ equal to 1 if n is a square and 0 otherwise, and let $r_2^+(n)$ be the number of representations of n as the form $i^2 + j^2$ where both i and j are positive integers. Then for $n \geq 1$,*

$$r_2^+(n) = \frac{1}{4} r_2(n) - \chi(n).$$

Proof. If n is not a square, then among the four representations $n = (\pm x)^2 + (\pm y)^2$ $(xy \neq 0)$ counted by $r_2(n)$, there is exactly one representation $|x|^2 + |y|^2$ counted by $r_2^+(n)$. Hence $r_2^+(n) = r_2(n)/4$.

If $n = k^2$ is a square, then we should first eliminate the four representations $k^2 + 0^2$, $(-k)^2 + 0^2$, $0^2 + k^2$, $0^2 + (-k)^2$ from the representations enumerated by $r_2(n)$. The result follows similarly by the arguments above. □

Proof of Theorem 2.1. Observe that $\varphi(q) = 1 + 2S(q)$ where $S(q) = \sum_{n\geq 1} q^{n^2}$. We have

$$\begin{aligned} \sum_{n=0}^{\infty} \overline{p}_k(n)(-q)^n &= \frac{1}{\varphi(q)^k} = \frac{1}{(1+2S(q))^k} \\ &\equiv \Big(1 - 2S(q) + 4S(q)^2 - 8S(q)^3\Big)^k \\ &\equiv 1 - 2kS(q) + 2k(k+1)S(q)^2 - \frac{4}{3}k(k+1)(k+2)S(q)^3 \pmod{16}. \end{aligned}$$

If $k \equiv 0, 3 \pmod 4$, then $\frac{4}{3}k(k+1)(k+2) \equiv 0 \pmod{16}$. If $k \equiv 1, 2 \pmod 4$, then $\frac{4}{3}k(k+1)(k+2) \equiv 0 \pmod 8$. We have

$$\sum_{n\geq 0} \overline{p}_k(n)(-q)^n \equiv 1 - 2kS(q) + 2k(k+1)S(q)^2 \pmod{2^{m(k)}}. \tag{2.1}$$

For $n \geq 1$, by the definition of $S(q)$ and $r_2^+(n)$ we obtain

$$(-1)^n \overline{p}_k(n) \equiv 2k(k+1) r_2^+(n) - 2k\chi(n) \pmod{2^{m(k)}}.$$

By Lemma 2.1 we can write

$$(-1)^n\overline{p}_k(n)\equiv\frac{1}{2}k(k+1)r_2(n)-2k(k+2)\chi(n)\pmod{2^{m(k)}}.\tag{2.2}$$

Let $n=2^\alpha(2n_0+1)$, $\alpha,n_0\geq 0$. From [2, Theorem 3.2.1] we see that

$$r_2(n)=4\sum_{\substack{d|n\\ d\,\text{odd}}}(-1)^{(d-1)/2}\equiv 4\sum_{\substack{d|n\\ d\,\text{odd}}}1\equiv 4d(2n_0+1)\pmod 8,$$

here $d(2n_0+1)$ denotes the number of positive divisors of $2n_0+1$. It is well-known that $d(2n_0+1)$ is odd if and only if $2n_0+1$ is a square. Hence we have

$$r_2(n)\equiv\begin{cases}4, & n\text{ is a square or twice a square}\\ 0, & \text{else}\end{cases}\pmod 8.\tag{2.3}$$

Note that if $k\equiv 0,3\pmod 4$, then $\frac{1}{2}k(k+1)\equiv 0\pmod 2$. Combining (2.2) with (2.3), we get the desired result. □

Since the number of integers not exceeding x which are a square or twice a square is less than $2\sqrt{x}$, we have

Corollary 2.1. *Given any integer $k\equiv 3\pmod 4$, $\overline{p}_k(n)$ is divisible by 16 for almost all positive integers n.*

In particular, for $k=3$, we have

$$\overline{p}_3(n)\equiv\begin{cases}10, & \text{if } n \text{ is an even square;}\\ 6, & \text{if } n \text{ is an odd square;}\\ 8, & \text{if } n \text{ is twice a square;}\\ 0, & \text{else}\end{cases}\pmod{16}.\tag{2.4}$$

This leads to the following corollary.

Corollary 2.2. *For any nonnegative integer n, we have*

$$\overline{p}_3(8n+1)\equiv 0\pmod 2,\quad \overline{p}_3(8n+2)\equiv 0\pmod 8,$$
$$\overline{p}_3(8n+3)\equiv 0\pmod{16},\quad \overline{p}_3(8n+4)\equiv 0\pmod 2,$$
$$\overline{p}_3(8n+5)\equiv 0\pmod{16},\quad \overline{p}_3(8n+6)\equiv 0\pmod{16}.$$

Proof. It follows directly from (2.4) that $\overline{p}_3(8n+1)\equiv\overline{p}_3(8n+4)\equiv 0\pmod 2$. Since any square is congruent to 0, 1 or 4 modulo 8, we know that $8n+2$ can not be a square. Hence by (2.4), we deduce that $\overline{p}_3(8n+2)\equiv 0\pmod 8$. Similarly, for $r\in\{3,5,6\}$, $8n+r$ cannot be a square or twice a square. Hence $\overline{p}_3(8n+r)\equiv 0\pmod{16}$. □

Since

$$\overline{p}_3(1)=6,\quad \overline{p}_3(2)=24,\quad \overline{p}_3(3)=80=2^4\cdot 5,\quad \overline{p}_3(4)=234=2\cdot 117,$$
$$\overline{p}_3(5)=624=2^4\cdot 39,\quad \overline{p}_3(6)=1552=2^4\cdot 97,$$

we see that all the modulus in this corollary cannot be replaced by higher powers of 2.

We have two deep Ramanujan-type congruences for $\overline{p}_3(n)$ modulo 32 and 64. This will be the next topic of this section. Before our discussion, we need some lemmas for preparation.

Lemma 2.2. *(Cf.* [13, Lemma 1]*.) The following identities hold:*

$$\varphi(q)=\varphi(q^4)+2q\psi(q^8),\quad \varphi(q)^2=\varphi(q^2)^2+4q\psi(q^4)^2,$$
$$\varphi(q)\varphi(-q)=\varphi(-q^2)^2,\quad \psi(q)^2=\varphi(q)\psi(q^2).$$

Lemma 2.3. *(Cf.* [3, Entry 25(viii)]*.)We have*

$$\varphi(q)^4-\varphi(-q)^4=16q\psi(q^2)^4.$$

Lemma 2.4. *(Cf.* [6, Lemma 2.1]*.) We have*

$$\frac{1}{\varphi(-q)}=\frac{1}{\varphi(-q^4)^4}\Big(\varphi(q^4)^3+2q\varphi(q^4)^2\psi(q^8)+4q^2\varphi(q^4)\psi(q^8)^2+8q^3\psi(q^8)^3\Big).$$

Theorem 2.2. *For any integer $n\geq 0$, we have*

$$\overline{p}_3(16n+14)\equiv 0 \pmod{32}.$$

Proof. Throughout this section, we will denote $a=\varphi(q^4)$ and $b=\psi(q^8)$. By Lemma 2.2, we have $\varphi(q)=a+2qb$. By Lemma 2.4, we have

$$\sum_{n\geq 0}\overline{p}_3(n)q^n=\frac{1}{\varphi(-q)^3}=\frac{1}{\varphi(-q^4)^{12}}\big(a^3+2qa^2b+4q^2ab^2+8q^3b^3\big)^3. \tag{2.5}$$

If we collect all the terms of the form q^{4n+2} in the expansion of $(a^3+2qa^2b+4q^2ab^2+8q^3b^3)^3$, we get

$$q^2(24a^7b^2+640a^3b^6q^4)\equiv 24q^2a^7b^2 \pmod{2^7}.$$

Collecting all the terms of the form q^{4n+2} on both sides of (2.5), dividing by q^2 and replacing q^4 by q, we obtain

$$\sum_{n\geq 0}\overline{p}_3(4n+2)q^n\equiv 24\frac{\varphi(q)^7\psi(q^2)^2}{\varphi(-q)^{12}} \pmod{2^7}. \tag{2.6}$$

By Lemma 2.3, we have $\varphi(-q)^4\equiv\varphi(q)^4 \pmod{2^4}$. Hence from (2.6) we deduce that

$$\sum_{n\geq 0}\overline{p}_3(4n+2)q^n\equiv 24\frac{\varphi(q)^7\psi(q^2)^2}{\varphi(q)^{12}}=24\frac{\psi(q^2)^2}{\varphi(q)^5} \pmod{2^7}. \tag{2.7}$$

Because $(a+2qb)^6\equiv a^6 \pmod 4$, we get $\varphi(q)^6\equiv\varphi(q^4)^6 \pmod 4$. Thus from (2.7) we obtain

$$\sum_{n\geq 0}\overline{p}_3(4n+2)q^n\equiv 24\frac{\psi(q^2)^2}{\varphi(q)^6}\varphi(q)\equiv 24\frac{\psi(q^2)^2}{\varphi(q^4)^6}\Big(\varphi(q^4)+2q\psi(q^8)\Big) \pmod{2^5}.$$

Collecting all the terms of the form q^{2n+1} on both sides, dividing by q and replacing q^2 by q, we obtain

$$\sum_{n\geq 0}\overline{p}_3(8n+6)q^n\equiv 16\frac{\psi(q)^2\psi(q^4)}{\varphi(q^2)^6}\equiv 16\frac{\psi(q^2)\psi(q^4)}{\varphi(q^2)^6} \pmod{32}.$$

Note that the terms of the form q^{2n+1} do not appear in the right hand side, we deduce that $\overline{p}_3\big(8(2n+1)+6\big)\equiv 0 \pmod{32}$, which is the desired result we want. □

Remark 2.1. The modulus 32 cannot be replaced by 64 since for $n = 0$, we have $\overline{p}_3(14) = 535008 = 2^5 \cdot 3 \cdot 5573$.

Theorem 2.3. *For any integer $n \geq 0$, we have*

$$\overline{p}_3(8n+7) \equiv 0 \pmod{64}.$$

Proof. We start from (2.5). If we collect all the terms of the form q^{4n+3} in the expansion of $(a^3 + 2qa^2b + 4q^2ab^2 + 8q^3b^3)^3$, we get

$$q^3(80a^6b^3 + 768a^2b^7q^4) \equiv 80q^3a^6b^3 \pmod{2^8}.$$

Collecting all the terms of the form q^{4n+3} on both sides of (2.5), then dividing by q^3 and replacing q^4 by q, we obtain

$$\sum_{n\geq 0} \overline{p}_3(4n+3)q^n \equiv 80\frac{\varphi(q)^6\psi(q^2)^3}{\varphi(-q)^{12}} \pmod{2^8}. \tag{2.8}$$

In the proof of Theorem 2.2 we have seen that $\varphi(q)^6 \equiv \varphi(q^4)^6 \pmod 4$. Hence $\varphi(-q)^6 \equiv \varphi(q^4)^6 \equiv \varphi(q)^6 \pmod 4$. We deduce from (2.8) that

$$\sum_{n\geq 0} \overline{p}_3(4n+3)q^n \equiv 80\frac{\psi(q^2)^3}{\varphi(q)^6} \equiv 80\frac{\psi(q^2)^3}{\varphi(q^4)^6} \pmod{2^6}.$$

Since the terms of the form q^{2n+1} do not appear in the right hand side, we deduce that $\overline{p}_3(4(2n+1)+3) \equiv 0 \pmod{64}$. This completes the proof. □

Remark 2.2. The modulus 64 in this theorem cannot be replaced by 128 as we have $\overline{p}_3(7) = 3648 = 2^6 \cdot 3 \cdot 19$.

3. Congruences Modulo 9

First we state two lemmas which will play a key role in our argument.

Lemma 3.1. *(Cf.* [1, Lemmas 2.6 and 2.7]*.) The following identities hold:*

$$\varphi(q) = \varphi(q^9) + 2qB(-q^3),$$

$$\varphi(q^3)^3 + 8qB(-q)^3 = \frac{\varphi(q)^4}{\varphi(q^3)},$$

where

$$B(q) = \frac{(q;q)_\infty (q^6;q^6)_\infty^2}{(q^2;q^2)_\infty (q^3;q^3)_\infty}.$$

Lemma 3.2. *(Cf.* [6, Lemma 2.1]*.) We have*

$$\frac{1}{\varphi(q)} = \frac{\varphi(q^9)}{\varphi(q^3)^4}\Big(\varphi(q^9)^2 - 2q\varphi(q^9)B(-q^3) + 4q^2B(-q^3)^2\Big).$$

The following lemma will also be used frequently.

Lemma 3.3. *Let p be a prime. We have*

$$(q;q)_\infty^p \equiv (q^p;q^p)_\infty \pmod p, \quad \varphi(q)^p \equiv \varphi(q^p) \pmod p.$$

Proof. By the binomial theorem, we have $(1-q)^p \equiv 1-q^p \pmod{p}$. This proves the first congruence. Since $\varphi(q) = (-q;-q)_\infty^2/(q^2;q^2)_\infty$, the second congruence relation follows from the first one. □

For convenience, we will let $s = \varphi(q^9)$ and $t = 2B(-q^3)$ throughout this section. From Lemmas 3.1 and 3.2, we have

$$s^3 + q^3t^3 = \frac{\varphi(q^3)^4}{\varphi(q^9)}, \tag{3.1}$$

$$\frac{1}{\varphi(q)} = \frac{\varphi(q^9)}{\varphi(q^3)^4}(s^2 - qst + q^2t^2). \tag{3.2}$$

We are going to establish an infinite family of congruences modulo $9 \cdot 2^4$.

Theorem 3.1. *We have*

$$\sum_{n\geq 0} \overline{p}_3(3n)(-q)^n \equiv \frac{\varphi(q^3)}{\varphi(q)^4} \pmod{9\cdot 2^3},$$

$$\sum_{n\geq 0} \overline{p}_3(3n+1)(-q)^n \equiv 6\frac{\varphi(q^3)^4}{\varphi(q)^8}B(-q) \pmod{9\cdot 2^4},$$

$$\sum_{n\geq 0} \overline{p}_3(3n+2)(-q)^n \equiv 24\frac{\varphi(q^3)^3}{\varphi(q)^8}B(-q)^2 \pmod{9\cdot 2^5}.$$

Proof. By (3.2) we have

$$\sum_{n\geq 0} \overline{p}_3(n)(-q)^n = \frac{1}{\varphi(q)^3} = \frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}(s^2 - qst + q^2t^2)^3. \tag{3.3}$$

Note that

$$(s^2 - qst + q^2t^2)^3 = (s^6 - 7q^3s^3t^3 + q^6t^6) + q(-3s^5t + 6q^3s^2t^4) + q^2(6s^4t^2 - 3q^3st^5).$$

Since $t = 2B(-q^3)$, collecting all the terms of the form q^{3n} on both sides of (3.3) and applying (3.1), we obtain

$$\begin{aligned}\sum_{n\geq 0} \overline{p}_3(3n)(-q)^{3n} &= \frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}(s^6 - 7q^3s^3t^3 + q^6t^6)\\ &\equiv \frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}(s^3 + q^3t^3)^2\\ &\equiv \frac{\varphi(q^9)}{\varphi(q^3)^4} \pmod{9\cdot 2^3}.\end{aligned}$$

Replacing q^3 by q, we get the first congruence identity.

Similarly, we have

$$\begin{aligned}\sum_{n\geq 0}\overline{p}_3(3n+1)(-q)^{3n+1} &= q\frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}(-3s^5t+6q^3s^2t^4)\\ &\equiv -3q\frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}s^2t(s^3+q^3t^3)\\ &\equiv -6q\frac{\varphi(q^9)^4}{\varphi(q^3)^8}B(-q^3) \pmod{9\cdot 2^4}.\end{aligned}$$

Dividing both sides by $-q$ and replacing q^3 by q, we get the second congruence identity.

In the same way, we have

$$\begin{aligned}\sum_{n\geq 0}\overline{p}_3(3n+2)(-q)^{3n+2} &= q^2\frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}(6s^4t^2-3st^5q^3)\\ &\equiv 6q^2\frac{\varphi(q^9)^3}{\varphi(q^3)^{12}}st^2(s^3+q^3t^3)\\ &\equiv 24q^2\frac{\varphi(q^9)^3}{\varphi(q^3)^8}B(-q^3)^2 \pmod{9\cdot 2^5}.\end{aligned}$$

Dividing both sides by q^2 and replacing q^3 by q, we get the third congruence identity. □

The following corollary follows immediately.

Corollary 3.1. *For any nonnegative integer n, we have*

$$\overline{p}_3(3n+1)\equiv 0 \pmod 6, \quad \overline{p}_3(3n+2)\equiv 0 \pmod{24}.$$

Theorem 3.2. *For any integer $\alpha\geq 1$, we have*

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha}n)(-q)^n \equiv \frac{\varphi(q^3)^4}{\varphi(q)^7} \pmod{9\cdot 2^3},$$

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha+1}n)(-q)^n \equiv \frac{\varphi(q^3)^5}{\varphi(q)^8} \pmod{9\cdot 2^3}.$$

Proof. We proceed our proof by induction on α.

By (3.2) and the first identity in Theorem 3.1, we have

$$\sum_{n\geq 0}\overline{p}_3(3n)(-q)^n \equiv \frac{\varphi(q^3)}{\varphi(q)^4} = \frac{\varphi(q^9)^4}{\varphi(q^3)^{15}}(s^2-qst+q^2t^2)^4 \pmod{9\cdot 2^3}.$$

Collecting all the terms of the form q^{3n} in the expansion of $(s^2-qst+q^2t^2)^4$ and reducing modulo 9, we get

$$s^8-16q^3s^5t^3+10q^6s^2t^6 \equiv s^2(s^3+q^3t^3)^2 \pmod{9\cdot 2^3}.$$

Hence we have

$$\sum_{n\geq 0}\overline{p}_3(9n)(-q)^{3n}\equiv\frac{\varphi(q^9)^4}{\varphi(q^3)^{15}}s^2(s^3+q^3t^3)^2=\frac{\varphi(q^9)^4}{\varphi(q^3)^7}\pmod{9\cdot 2^3}.$$

Replacing q^3 by q we proved the first identity for $\alpha=1$.

Suppose

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha}n)(-q)^n\equiv\frac{\varphi(q^3)^4}{\varphi(q)^7}\pmod{9\cdot 2^3}. \tag{3.4}$$

Then by (3.2) we have

$$\frac{\varphi(q^3)^4}{\varphi(q)^7}=\frac{\varphi(q^9)^7}{\varphi(q^3)^{24}}(s^2-qst+q^2t^2)^7.$$

Collecting those terms of the form q^{3n} in the expansion of $(s^2-qst+q^2t^2)^7$ and reducing modulo 9, we get

$$\begin{aligned}&s^{14}-77q^3s^{11}t^3+357q^6s^8t^6-266q^9s^5t^9+28q^{12}s^2t^{12}\\&\equiv s^{14}+4q^3s^{11}t^3+6q^6s^8t^6+4q^9s^5t^9+q^{12}s^2t^{12}\\&=s^2(s^3+q^3t^3)^4\pmod{9\cdot 2^3}.\end{aligned}$$

If we collect all the terms of the form q^{3n} in (3.4), we get

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha+1}n)(-q)^{3n}\equiv\frac{\varphi(q^9)^7}{\varphi(q^3)^{24}}s^2(s^3+q^3t^3)^4\equiv\frac{\varphi(q^9)^5}{\varphi(q^3)^8}\pmod{9\cdot 2^3}.$$

Replacing q^3 by q, we obtain

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha+1}n)(-q)^n\equiv\frac{\varphi(q^3)^5}{\varphi(q)^8}\pmod{9\cdot 2^3}. \tag{3.5}$$

Now suppose we have (3.5), by (3.2) we deduce that

$$\frac{\varphi(q^3)^5}{\varphi(q)^8}=\frac{\varphi(q^9)^8}{\varphi(q^3)^{27}}(s^2-qst+q^2t^2)^8.$$

Collecting all the terms of the form q^{3n} in the expansion of $(s^2-qst+q^2t^2)^8$ and reducing modulo 9, we get

$$\begin{aligned}&s^{16}-112q^3s^{13}t^3+784q^6s^{10}t^6-1016q^9s^7t^9+266q^{12}s^4t^{12}-8q^{15}st^{15}\\&\equiv s(s^{15}+5q^3s^{12}t^3+10q^6s^9t^6+10q^9s^6t^9+5q^{12}s^3t^{12}+q^{15}t^{15})\\&=s(s^3+q^3t^3)^5\pmod{9\cdot 2^3}.\end{aligned}$$

If we collect all the terms of the form q^{3n} in (3.5), we get

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha+2}n)(-q)^{3n}\equiv\frac{\varphi(q^9)^8}{\varphi(q^3)^{27}}s(s^3+q^3t^3)^5\equiv\frac{\varphi(q^9)^4}{\varphi(q^3)^7}\pmod{9\cdot 2^3}.$$

Replacing q^3 by q, we obtain

$$\sum_{n\geq 0}\overline{p}_3(3^{2\alpha+2}n)(-q)^n\equiv\frac{\varphi(q^3)^4}{\varphi(q)^7}\pmod{9\cdot 2^3}.$$

By inducting on α, we complete our proof. □

Theorem 3.3. *For any integers $\alpha\geq 1$ and $n\geq 0$, we have*

$$\overline{p}_3\big(3^{2\alpha+1}(3n+2)\big)\equiv 0\pmod{144}.$$

Proof. By (3.2) we have

$$\frac{\varphi(q^3)^5}{\varphi(q)^8}=\frac{\varphi(q^9)^8}{\varphi(q^3)^{27}}(s^2-qst+q^2t^2)^8.$$

If we collect all the terms of the form q^{3n+2} in the expansion of $(s^2-qst+q^2t^2)^8$, we get

$$9q^2(4s^{14}t^2-56q^3s^{11}t^5+123q^6s^8t^8-56q^9s^5t^{11}+4q^{12}s^2t^{14}).$$

It is obvious that all the coefficients are divisible by 9. Thus all the coefficients of the terms q^{3n+2} $(n=0,1,2,\cdots)$ in both sides of (3.5) are divisible by 9, which implies $\overline{p}_3(3^{2\alpha+1}(3n+2))\equiv 0\pmod 9$.

For any integer x, we have $x^2\equiv 0,1,4,7\pmod 9$. Hence $3^{2\alpha+1}(3n+2)=9^\alpha(9n+6)$ cannot be a square. Moreover, since $2x^2\equiv 0,2,5,8\pmod 9$, we know $9n+6$ cannot be twice a square. Thus $9^\alpha(9n+6)$ cannot be twice a square. By (2.4) we know $\overline{p}_3(3^{2\alpha+1}(3n+2))\equiv 0\pmod{16}$. Combining this with $\overline{p}_3(3^{2\alpha+1}(3n+2))\equiv 0\pmod 9$, we complete our proof. □

Corollary 3.2. *$\overline{p}_3(n)$ is divisible by 144 for at least $\frac{1}{72}$ of the time.*

Proof. The arithmetic sequences $\big\{3^{2\alpha+1}(3n+2):n\geq 0\big\}$ $(\alpha=1,2,\cdots)$ mentioned in Theorem 3.3 are disjoint for different α, and they account for

$$\frac{1}{3^4}+\frac{1}{3^6}+\cdots=\frac{1}{72}$$

of all positive integers. □

4. Congruences Modulo 7 and 11

Theorem 4.1. *For any integer $n\geq 1$, we have*

$$\overline{p}_3(7n)\equiv(-1)^n r_3(n)\pmod 7.$$

To prove this theorem, we need the following lemma.

Lemma 4.1. *(Cf.* [21, Lemma 12]*.) For any prime $p\geq 3$, we have*

$$r_4(pn)\equiv r_4(n)\pmod p,\quad r_8(pn)\equiv r_8(n)\pmod p.$$

Proof of Theorem 4.1. We have

$$\varphi(q)^7\sum_{n\geq 0}\overline{p}_3(n)(-q)^n=\varphi(q)^4=\sum_{n\geq 0}r_4(n)q^n.$$

By Lemma 3.3, we have $\varphi(q)^7\equiv\varphi(q^7)\pmod 7$. Hence

$$\varphi(q^7)\sum_{n\geq 0}\overline{p}_3(n)(-q)^n\equiv\sum_{n\geq 0}r_4(n)q^n\pmod 7.$$

Collecting all the terms of the form q^{7n} on both sides of this identity, then replacing q^7 by q and applying Lemma 4.1 with $p=7$, we obtain

$$\varphi(q)\sum_{n\geq 0}\overline{p}_3(7n)(-q)^n \equiv \sum_{n\geq 0} r_4(7n)q^n \equiv \sum_{n\geq 0} r_4(n)q^n = \varphi(q)^4 \pmod 7.$$

This implies

$$\sum_{n\geq 0}\overline{p}_3(7n)(-q)^n \equiv \varphi(q)^3 = \sum_{n\geq 0} r_3(n)q^n \pmod 7. \tag{4.1}$$

The theorem follows by comparing the coefficients of q^n on both sides. □

Theorem 4.2. *For any nonnegative integers α and n, we have*

$$\overline{p}_3(7\cdot 4^{\alpha+1}n) \equiv (-1)^n\overline{p}_3(7n) \pmod 7$$

and

$$\overline{p}_3(4^\alpha(56n+49)) \equiv 0 \pmod 7.$$

Proof. From (4.1) and Lemma 2.2 we see that

$$\begin{aligned}\sum_{n=0}^{\infty}\overline{p}_3(7n)q^n &\equiv \varphi(-q)^3 = (\varphi(q^4)-2q\psi(q^8))^3\\ &= \varphi(q^4)^3 - 6q\varphi(q^4)^2\psi(q^8) + 12q^2\psi(q^8)^2\varphi(q^4) - 8q^3\psi(q^8)^3 \pmod 7.\end{aligned} \tag{4.2}$$

Extracting the terms of the form q^{4n} on both sides, replacing q^4 by q, we obtain

$$\sum_{n=0}^{\infty}\overline{p}_3(7\cdot 4n)q^n \equiv \varphi(q)^3 = \sum_{n=0}^{\infty}(-1)^n\overline{p}_3(7n)q^n \pmod 7.$$

Hence we have $\overline{p}_3(7\cdot 4n) \equiv (-1)^n\overline{p}_3(7n) \pmod 7$, from which the first congruence follows.

Extracting the terms of the form q^{4n+3} on both sides of (4.2), dividing by q^3 and replacing q^4 by q, we obtain

$$\sum_{n=0}^{\infty}\overline{p}_3(7\cdot(4n+3))q^n \equiv -8\psi(q^2)^3 \pmod 7.$$

Since the terms of the form q^{2n+1} do not appear on the right hand side, we deduce that $\overline{p}_3(7(8n+7)) \equiv 0 \pmod 7$. Combining this congruence with the relation $\overline{p}_3(7\cdot 4^{\alpha+1}n) \equiv (-1)^n\overline{p}_3(7n) \pmod 7$, we proved the second congruence. □

Theorem 4.3. *Let $p\geq 3$ be a prime, and N be a positive integer which is coprime to p. Let α be any nonnegative integer.*
(1) If $p\equiv 1 \pmod 7$, then $\overline{p}_3(7p^{14\alpha+13}N)\equiv 0 \pmod 7$.
(2) If $p\equiv 2,3,4,5,6 \pmod 7$, then $\overline{p}_3(7p^{12\alpha+11}N)\equiv 0 \pmod 7$.
(3) If $p\equiv 6 \pmod 7$, then $\overline{p}_3(7p^3N)\equiv 0 \pmod 7$.

Proof. From [11, (4)] we have

$$r_3(p^2n) + \Big(\frac{-n}{p}\Big)r_3(n) + pr_3(n/p^2) = (p+1)r_3(n), \tag{4.3}$$

here $(\frac{\cdot}{p})$ is the Legendre symbol and we take $r_3(n/p^2)=0$ unless $p^2|n$. Replacing n by np, we see that

$$r_3(p^3n) + pr_3(n/p) = (p+1)r_3(pn).$$

By Theorem 4.1, we have

$$\overline{p}_3(7p^3n)+p\overline{p}_3(7n/p)\equiv(p+1)\overline{p}_3(7pn)\pmod 7. \tag{4.4}$$

From (4.4), by iterating we obtain

$$\overline{p}_3(7p^{2k+1}n)\equiv\frac{p(p^k-1)}{p-1}\Big(\overline{p}_3(7pn)-\overline{p}_3\big(7n/p\big)\Big)+\overline{p}_3(7pn)\pmod 7.$$

Let $n=N$, which is coprime to p. We get $\overline{p}_3(7n/p)=0$ and thus

$$\overline{p}_3(7p^{2k+1}N)\equiv\Big(\frac{p(p^k-1)}{p-1}+1\Big)\overline{p}_3(7pN)\pmod 7. \tag{4.5}$$

(1) Let $k=7\alpha+6$ in (4.5). Since $p\equiv 1 \pmod 7$, we have

$$\frac{p^{7\alpha+6}-1}{p-1}=1+p+\cdots+p^{7\alpha+5}\equiv -1\pmod 7.$$

Hence by (4.5) we deduce that $\overline{p}_3(7p^{14\alpha+13}N)\equiv 0 \pmod 7$.

(2) Let $k=6\alpha+5$ in (4.5). In this case, we have $p^{6\alpha+6}\equiv 1 \pmod 7$. Hence from (4.5) we obtain

$$\overline{p}_3(7p^{12\alpha+11}N)\equiv\Big(\frac{p^{6\alpha+6}-p}{p-1}+1\Big)\overline{p}_3(7pN)\equiv 0\pmod 7.$$

(3) Let $n=N$ in (4.4). Since $p+1\equiv 0 \pmod 7$, the congruence follows immediately. □

Theorem 4.4. *For any integers $n\geq 0$ and $\alpha\geq 1$, we have*

$$\overline{p}_3(7^4\cdot n)\equiv\overline{p}_3(7^2\cdot n)\pmod 7$$

and

$$\overline{p}_3\big(7^{2\alpha+1}(7n+3)\big)\equiv\overline{p}_3\big(7^{2\alpha+1}(7n+5)\big)\equiv\overline{p}_3\big(7^{2\alpha+1}(7n+6)\big)\equiv 0\pmod 7.$$

Proof. By Theorem 4.1 and (4.3) we obtain

$$\overline{p}_3(7\cdot p^2n)+\Big(\frac{-n}{p}\Big)\overline{p}_3(7n)+p\overline{p}_3(7n/p^2)\equiv(p+1)\overline{p}_3(7n)\pmod 7. \tag{4.6}$$

Let $p=7$ and we replace n by $7n$. We deduce that $\overline{p}_3(7^4\cdot n)\equiv\overline{p}_3(7^2\cdot n) \pmod 7$.

Let $p=7$ and $n=7m+r, r\in\{3,5,6\}$ in (4.6). Note that $\big(\frac{-r}{7}\big)=1$, it follows from (4.6) that $\overline{p}_3(7^3(7m+r))\equiv 0 \pmod 7$. Combining this congruence with the first congruence relation, by induction on α, we complete our proof of the theorem. □

Corollary 4.1. *$\overline{p}_3(n)$ is divisible by 7 for at least $1/784$ of all the time.*

Proof. Note that for different pairs (α,r), where $\alpha\geq 1$ and $r\in\{3,5,6\}$, the arithmetic sequences $\big\{7^{2\alpha+1}(7n+r):n=0,1,2,\cdots\big\}$ are disjoint, and they account for

$$3\Big(\frac{1}{7^4}+\frac{1}{7^6}+\cdots+\frac{1}{7^{2\alpha+2}}+\cdots\Big)=\frac{1}{784}$$

of all nonnegative integers. □

Theorem 4.5. *For any integers $n\geq 0$, we have*

$$\overline{p}_3(11n)\equiv(-1)^n r_7(n)\pmod{11}.$$

Proof. By definition we have

$$\varphi(q)^{11}\sum_{n\geq 0}\overline{p}_3(n)(-q)^n=\varphi(q)^8=\sum_{n\geq 0}r_8(n)q^n.$$

By Lemma 3.3, we have $\varphi(q)^{11}\equiv\varphi(q^{11})\pmod{11}$. Hence

$$\varphi(q^{11})\sum_{n\geq 0}\overline{p}_3(n)(-q)^n\equiv\sum_{n\geq 0}r_8(n)q^n\pmod{11}.$$

Collecting all the terms of the form q^{11n} on both sides, then replacing q^{11} by q and applying Lemma 4.1 with $p=11$, we get

$$\varphi(q)\sum_{n\geq 0}\overline{p}_3(11n)(-q)^n\equiv\sum_{n\geq 0}r_8(11n)q^n\equiv\sum_{n\geq 0}r_8(n)q^n=\varphi(q)^8\pmod{11}.$$

Hence

$$\sum_{n\geq 0}\overline{p}_3(11n)(-q)^n\equiv\varphi(q)^7=\sum_{n\geq 0}r_7(n)q^n\pmod{11}.$$

The theorem follows by comparing the coefficients of q^n on both sides. □

Theorem 4.6. *Let $p\geq 3$ be a prime, and N be a positive integer which is coprime to p. Let α be any nonnegative integer.*
(1) We have $\overline{p}_3(11^3\cdot n)\equiv\overline{p}_3(11n)\pmod{11}$.
(2) If $p\equiv 1,3,4,5,9\pmod{11}$, then $\overline{p}_3(11p^{22\alpha+21}N)\equiv 0\pmod{11}$.
(3) If $p\equiv 2,6,7,8,10\pmod{11}$, then $\overline{p}_3(11p^{4\alpha+3}N)\equiv 0\pmod{11}$.

Proof. From [9, Lemma 5.1] we have

$$r_7(p^2n)=\Big(p^5-p^2\Big(\frac{-n}{p}\Big)+1\Big)r_7(n)-p^5r_7(n/p^2),\tag{4.7}$$

here as before, $(\frac{\cdot}{p})$ is the Legendre symbol and we take $r_7(n/p^2)=0$ unless $p^2|n$. Replacing n by np, we see that

$$r_7(p^3n)=(p^5+1)r_7(pn)-p^5r_7(n/p).$$

Hence by Theorem 4.5 we have

$$\overline{p}_3(11p^3n)\equiv(p^5+1)\overline{p}_3(11pn)-p^5\overline{p}_3(11n/p)\pmod{11}.\tag{4.8}$$

From (4.8), by iterating we see that

$$\overline{p}_3(11p^{2k+1}n)\equiv\frac{p^5(p^{5k}-1)}{p^5-1}\Big(\overline{p}_3(11pn)-\overline{p}_3(11n/p)\Big)+\overline{p}_3(11pn)\pmod{11}.$$

Let $n=N$, which is coprime to p. We get $\overline{p}_3(11n/p)=0$ and thus

$$\overline{p}_3(11p^{2k+1}N)\equiv\Big(\frac{p^5(p^{5k}-1)}{p^5-1}+1\Big)\overline{p}_3(11pN)\pmod{11}.\tag{4.9}$$

(1) Let $p=11$ in (4.7). We deduce that $r_7(11^2\cdot n)\equiv r_7(n)\pmod{11}$. By Theorem 4.5 we get the desired congruence we want.

(2) Let $k=11\alpha+10$ in (4.9). Note that $p\equiv 1,3,4,5,9\pmod{11}$ implies $p^5\equiv 1\pmod{11}$. We have

$$\frac{p^{55\alpha+50}-1}{p^5-1}=1+p^5+p^{10}+\cdots+p^{5(11\alpha+9)}\equiv 10\pmod{11}.$$

It follows from (4.5) that $\overline{p}_3(11p^{22\alpha+21}N) \equiv 0 \pmod{11}$.

(3) Let $k = 2\alpha + 1$ in (4.9). Note that $p \equiv 2, 6, 7, 8, 10 \pmod{11}$ implies $p^5 \equiv -1 \pmod{11}$. It follows from (4.9) that $\overline{p}_3(11p^{4\alpha+3}N) \equiv 0 \pmod{11}$. □

For example, let $p = 7$ and $N = 7n+r$, $r \in \{1, 2, 3, 4, 5, 6\}$ in part (2) of Theorem 4.6, we have

$$\overline{p}_3\big(11 \cdot 7^{4\alpha+3}(7n + r)\big) \equiv 0 \pmod{11}.$$

By a similar argument to the proof of Corollary 4.1, we can prove

Corollary 4.2. *$\overline{p}_3(n)$ is divisible by 11 for at least $1/4400$ of all the time.*

References

1. G. E. Andrews, M. D. Hirschhorn, and J. A. Sellers, Arithmetic properties of partitions with even parts distinct, Ramanujan J. **23** (2010), 169–181.
2. B. C. Berndt, Number Theory in the Spirit of Ramanujan, Am. Math. Soc., Providence, 2006.
3. B. C. Berndt, Ramanujan's Notebooks, Part III, Springer-Verlag, New York, 1991.
4. K. Bringmann and J. Lovejoy, Rank and congruences for overpartition pairs, Int. J. Number Theory **4** (2008), 303–322.
5. S. C. Chen, Congruences for overpartition k-tuples, Integers **10** (2010), 477–484.
6. W. Y. C. Chen and B. L. S. Lin, Arithmetic properties of overpartition pairs, Acta Arith. **151** (2012), 263–277.
7. W. Y. C. Chen, L. H. Sun, R.-H. Wang, and L. Zhang, Ramanujan-type congruences for overpartitions modulo 5, preprint, arXiv:1406.3801.
8. W. Y. C. Chen and E. X. W. Xia, Proof of a conjecture of Hirschhorn and Sellers on overpartitions, Acta Arith. **163** (2014), 59–69.
9. S. Cooper, Sums of five, seven and nine squares, Ramanujan J. **6** (2002), 469–490.
10. S. Corteel and J. Lovejoy, Overpartitions, Trans. Amer. Math. Soc. **356** (2004), 1623–1635.
11. M. D. Hirschhorn and J. A. Sellers, On representations of a number as a sum of three squares, Discrete Math. **199** (1999), 85–101.
12. M. D. Hirschhorn and J. A. Sellers, An infinite family of overpartition congruences modulo 12, Integers **5** (2005), Article A20.
13. M. D. Hirschhorn and J. A. Sellers, Arithmetic relations for overpartitions, J. Combin. Math. Combin. Comput. **53** (2005), 65–73.
14. B. Kim, The overpartition function modulo 128, Integers **8** (2008), Article A38.
15. B. Kim, A short note on the overpartition function, Discrete Math. **309** (2009), 2528–2532.
16. B. Kim, Overpartition pairs modulo powers of 2, Discrete Math. **311** (2011), 835–840.
17. B. L. S. Lin, A new proof of a conjecture of Hirschhorn and Sellers on overpartitions, Ramanujan J, doi:10.1007/s11139-014-9598-z, to appear.
18. D. M. Keister, J. A. Sellers, and R. G. Vary, Some arithmetic properties of overpartition tuples, Integers **9** (2009), A17, 181–190.
19. J. Lovejoy, Overpartition theorems of the Rogers-Ramanujan type, J. London Math. Soc. **69** (2004), 562–574.
20. K. Mahlburg, The overpartition function modulo small powers of 2, Discrete. Math. **286** (3) (2004), 263–267.
21. L. Wang, Another proof of a conjecture of Hirschhorn and Sellers on overpartitions, J. Integer Seq. **17** (2014), Article 14.9.8.

Department of Mathematics, National University of Singapore, Singapore, 119076
E-mail address: wangliuquan@u.nus.edu; mathlqwang@163.com